
**Space environment (natural and
artificial) — Galactic cosmic ray model**

*Environnement spatial (naturel et artificiel) — Modèle de rayonnement
cosmique galactique*

Not Official Standard
For Review Only



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

Not Official Standard
For Review Only

© ISO 2004

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Licensed to: Tongson Nick Mr.
Downloaded: 2012-09-10
Single user licence only, copying and networking prohibited

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 15390 was prepared by Technical Committee ISO/TC 20, *Aircraft and space vehicles*, Subcommittee SC 14, *Space systems and operations*.

Not Official Standard
For Review Only

Not Official Standard
For Review Only

Space environment (natural and artificial) — Galactic cosmic ray model

1 Scope

This International Standard specifies a model for estimating the radiation impact of galactic cosmic rays (GCR) on hardware and on biological and other objects when in space. This International Standard can also be used in scientific research to generalize the available experimental evidence for GCR fluxes. This International Standard establishes the model parameters and characteristics of variations in the 10^1 MeV to 10^5 MeV GCR particles (electrons, protons, and $Z = 2$ to 92 nuclei in the near-Earth space beyond the Earth's magnetosphere).

2 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

2.1

galactic cosmic rays

GCR

high-energy charged particle fluxes penetrating the heliosphere from local interstellar space

2.2

Wolf number

W

$$W = 10g+f$$

where

g is sunspot group number;

f is the total sunspot number on the sun's visible disk

2.3

rigidity spectrum

$\Phi_i(R)$

rigidity distribution of cosmic ray particle fluxes

2.4

energy spectrum

$F_i(E)$

energy distribution of cosmic ray particle fluxes

3 Principles of the model

3.1 The model describes the variations of GCR fluxes due to variations in solar activity and in the large-scale heliospheric magnetic field (the sun's polar magnetic field) throughout 22-year cycles.

3.2 The angular distribution of galactic cosmic ray fluxes in the Earth's orbit beyond the Earth's magnetosphere is to be isotropic.

3.3 Solar activity is characterized by 12-month averages of Wolf numbers (sunspot numbers) \overline{W} .

3.4 The variations of the large-scale heliospheric magnetic field are assumed to be proportional to the variations of the Sun's polar magnetic field whose intensity and polarity are taken to be dependent on solar activity and on whether a given solar cycle is even or odd:

$$M[\overline{W}(t), n] = (-1)^{n-1} \times S \times \left\{ 1 - \left[\frac{\overline{W}(t) - \overline{W}_n^{\min}}{\overline{W}_n^{\max} - \overline{W}_n^{\min}} \right]^{2,7} \right\} \quad (1)$$

where

$$S = 1 \text{ at } t - t_n^{\pm} \geq 0 \text{ (otherwise, } S = -1);$$

t_n^{\pm} is the sign reversal moment of the polar magnetic field in the n th solar cycle identified with solar maximum (values suggested for particle flux prediction: $t_{19}^{\pm} = 1958,21$; $t_{20}^{\pm} = 1968,87$; $t_{21}^{\pm} = 1979,96$; $t_{22}^{\pm} = 1989,46$; $t_{23}^{\pm} = 2\,000,71$ and $t_{24}^{\pm} = 2\,011,3$);

W_n^{\max} is the highest level of solar activity in the n th solar cycle;

W_n^{\min} is the least solar activity level that borders the n th solar cycle.

3.5 The dynamics of the large-scale GCR modulation is characterized by the effective modulation potential of the heliosphere, $R_0(t, R)$, (for particles of rigidity R at a given moment t) calculated as

$$R_0 \left\{ \overline{W} [t - \Delta t(n, R, t)] \right\} = 0,37 + 3 \times 10^{-4} \times W^{1,45} [t - \Delta t(n, R, t)] \quad (2)$$

3.6 The lag, $\Delta t(n, R, t)$, of GCR flux variations relative to solar activity variations is taken to depend on magnetic rigidity, R , of particles, on whether a solar cycle is odd or even (n), and on solar cycle phase:

$$\Delta t(n, R, t) = 0,5 \times [T_+ + T_-(R)] + 0,5 \times [T_+ - T_-(R)] \times \tau(\overline{W}) \quad (3)$$

where the lag amplitude is independent of particle rigidity in even cycles (T_+):

$$T_+ = 15 \text{ months} \quad (4)$$

and in odd cycles (T_-) is

$$T_-(R) = 7,5 \times R^{-0,45} \text{ months} \quad (5)$$

The time function of the lag variations from Equation (3) to be

$$\tau(\overline{W}) = (-1)^n \times \left[\frac{\overline{W}(t - \delta_w t) - \overline{W}_n^{\min}}{\overline{W}_n^{\max}} \right]^{0,2} \quad (6)$$

where $\delta_w t = 16$ months.

4 GCR rigidity and energy spectra

4.1 General

In terms of the GCR model, the particle flux rigidity and energy spectra are calculated consecutively.

4.2 GCR particle rigidity spectra

4.2.1 GCR particle rigidity spectra $\Phi_i(R, t)$ ($\text{s}\cdot\text{m}^2\cdot\text{sr}\cdot\text{GeV}^{-1}$) for particles of rigidity R at moment t are calculated using the following equation:

$$\Phi_i(R, t) = \frac{C_i \times \beta^{\alpha_i}}{R^{\gamma_i}} \times \left[\frac{R}{R + R_0(R, t)} \right]^{\Delta_i(R, t)} \quad (7)$$

where

$\Delta_i(R, t)$ is a dimensionless parameter calculated using the following equation:

$$\Delta_i(R, t) = 5,5 + 1,13 \frac{Z_i}{|Z_i|} M[\bar{W}(t), n] \times \frac{\beta \times R}{R_0(R, t)} \exp\left[\frac{\beta \times R}{R_0(R, t)} \right] \quad (8)$$

where β is the particle velocity-to-luminal velocity ratio:

$$\beta = \frac{R}{\sqrt{R^2 + \left(\frac{A_i m_i}{|Z_i|} \right)^2}} \quad (9)$$

where A_i and Z_i are particle mass number and particle charge respectively (see Tables 1 and 2);

m_i is particle rest mass, namely,

$$m_e = 5,1 \times 10^{-4} \text{ GeV for electrons,}$$

$$m_p = 0,938 \text{ GeV for protons,}$$

$$m_{Z \geq 2} = 0,939 \text{ GeV/nucleon for nuclei.}$$

C_i, α_i, γ_i are parameters of non-modulated rigidity spectrum of i -specie particles. For particles with $Z \leq 28$ see Table 1. For particles with $Z \geq 29$, $\alpha_i = \alpha_{26}$ and $\gamma_i = \gamma_{26}$, C_i is calculated from equation

$$C_i = C_{26} \frac{C_i}{C_{26}}$$

where $\frac{C_i}{C_{26}}$ data is found in Table 2.

4.2.2 The standard deviation values, $[\sigma_{\Phi_i(R,t)}]$, are calculated using the following equation:

$$\sigma_{\Phi_i(R,t)} = \Phi_i(R,t) \times \sqrt{\left(\frac{\sigma_{C_i}}{C_i}\right)^2 + \frac{0,08}{\left[1 + \frac{R}{R_0(R,t)}\right]^2}} \quad (10)$$

4.3 GCR energy spectra

4.3.1 The energy spectra $F_i(E,t)$ ($\text{s}\cdot\text{m}^2\cdot\text{sr}\cdot\text{GeV}^{-1}$) of GCR particles of energy E at moment t are calculated using the following equation:

$$F_i(E,t) = \Phi_i(R,t) \frac{A_i}{|Z_i|} \frac{10^{-3}}{\beta} \quad (11)$$

For particles with $A_i = 1$, the units are $(\text{s}\cdot\text{m}^2\cdot\text{sr}\cdot\text{MeV})^{-1}$.

For particles with $A_i \geq 2$, the units are $(\text{s}\cdot\text{m}^2\cdot\text{sr}\cdot\text{MeV}/\text{nucleon})^{-1}$.

4.3.2 With prescribed rigidity R of GCR particles, the kinetic energy E , in GeV (GeV/nucleon for nuclei), is calculated using the following equation:

$$E = -m_i + \sqrt{m_i^2 + \left(\frac{Z_i}{A_i} R\right)^2} \quad (12)$$

4.3.3 With prescribed kinetic energy of particles, the rigidity R and the relative velocity β of the particles are calculated using the following equation:

$$R = \frac{A_i}{|Z_i|} \sqrt{E(E + 2m_i)} \quad (13)$$

$$\beta = \frac{\sqrt{E(E + 2m_i)}}{E + m_i} \quad (14)$$

4.3.4 Standard deviation values $[\sigma_{F_i(E,t)}]$ are calculated using the following equation:

$$\sigma_{F_i(E,t)} = \frac{\sigma_{\Phi_i(R,t)}}{\Phi_i(R,t)} \times F_i(E,t) \quad (15)$$

Table 1 — Parameters of GCR rigidity spectra for particles $Z \leq 28$

Z	Particle	A_i	$C_i \pm \sigma_{C_i}$	γ_i	α_i
-1	e	1,0	170	γ_e^a	—
1	H	1,0	$(1,85 \pm 0,13) 10^4$	2,74 0,02	2,85 0,02
2	He	4,0	$(3,69 \pm 0,22) 10^3$	$2,77 \pm 0,02$	$3,12 \pm 0,02$
3	Li	6,9	$19,5 \pm 1,5$	$2,82 \pm 0,02$	$3,41 \pm 0,11$
4	Be	9,0	$17,7 \pm 1,3$	$3,05 \pm 0,02$	$4,30 \pm 0,12$
5	B	10,8	$49,2 \pm 1,6$	$2,96 \pm 0,01$	$3,93 \pm 0,05$
6	C	12,0	$103,0 \pm 3,0$	$2,76 \pm 0,01$	$3,18 \pm 0,04$
7	N	14,0	$36,7 \pm 1,2$	$2,89 \pm 0,01$	$3,77 \pm 0,05$
8	O	16,0	$87,4 \pm 2,1$	$2,70 \pm 0,01$	$3,11 \pm 0,04$
9	F	19,0	$3,19 \pm 0,28$	$2,82 \pm 0,03$	$4,05 \pm 0,06$
10	Ne	20,2	$16,4 \pm 0,70$	$2,76 \pm 0,01$	$3,11 \pm 0,07$
11	Na	23,0	$4,43 \pm 0,28$	$2,84 \pm 0,02$	$3,14 \pm 0,09$
12	Mg	24,3	$19,3 \pm 0,70$	$2,70 \pm 0,01$	$3,65 \pm 0,27$
13	Al	27,0	$4,17 \pm 0,22$	$2,77 \pm 0,02$	$3,46 \pm 0,21$
14	Si	28,1	$13,4 \pm 0,50$	$2,66 \pm 0,01$	$3,00 \pm 0,10$
15	P	31,0	$1,15 \pm 0,04$	$2,89 \pm 0,01$	$4,04 \pm 0,41$
16	S	32,1	$3,06 \pm 0,12$	$2,71 \pm 0,02$	$3,30 \pm 0,22$
17	Cl	35,4	$1,30 \pm 0,08$	$3,00 \pm 0,04$	$4,40 \pm 0,30$
18	Ar	39,9	$2,33 \pm 0,07$	$2,93 \pm 0,01$	$4,33 \pm 0,21$
19	K	39,1	$1,87 \pm 0,05$	$3,05 \pm 0,01$	$4,49 \pm 0,20$
20	Ca	40,1	$2,17 \pm 0,06$	$2,77 \pm 0,01$	$2,93 \pm 0,16$
21	Sc	44,9	$0,74 \pm 0,02$	$2,97 \pm 0,01$	$3,78 \pm 0,19$
22	Ti	47,9	$2,63 \pm 0,08$	$2,99 \pm 0,01$	$3,79 \pm 0,17$
23	V	50,9	$1,23 \pm 0,04$	$2,94 \pm 0,01$	$3,50 \pm 0,14$
24	Cr	52,0	$2,12 \pm 0,06$	$2,89 \pm 0,01$	$3,28 \pm 0,17$
25	Mn	54,9	$1,14 \pm 0,05$	$2,74 \pm 0,02$	$3,29 \pm 0,27$
26	Fe	55,8	$9,32 \pm 0,24$	$2,63 \pm 0,01$	$3,01 \pm 0,07$
27	Co	58,9	$0,10 \pm 0,08$	2,63	$4,25 \pm 0,79$
28	Ni	58,7	$0,49 \pm 0,02$	$2,63 \pm 0,01$	$3,52 \pm 0,28$

NOTE 1 In the case of $Z > 2$ nuclei, the C_i , γ_i , α_i values are for a mixture of the respective isotopes.

NOTE 2 Atomic masses corresponding to the natural elemental abundances are taken to be mass numbers A_i for $Z > 2$ nuclei in conformity with the periodic system chart. This is within the model accuracy.

^a In the case of electrons, the parameter $\gamma_e = 3,0 - 1,4 \exp\left(\frac{R}{R_e}\right)$, where $R_e = 1$ GeV.

Table 2 — Parameters and GCR rigidity spectra for particles $Z \geq 29$

Z	Particle	A_i	C_i/C_{26}
29	Cu	63,5	$6,8 \times 10^{-4}$
30	Zn	65,4	$8,8 \times 10^{-4}$
31	Ga	69,7	$6,5 \times 10^{-5}$
32	Ge	72,6	$1,4 \times 10^{-4}$
33	As	74,9	$8,9 \times 10^{-6}$
34	Se	79,0	$5,2 \times 10^{-5}$
35	Br	79,9	$9,7 \times 10^{-6}$
36	Kr	83,8	$2,7 \times 10^{-5}$
37	Rb	85,5	$8,8 \times 10^{-6}$
38	Sr	87,6	$2,9 \times 10^{-5}$
39	Y	88,9	$6,5 \times 10^{-6}$
40	Zr	91,2	$1,6 \times 10^{-5}$
41	Nb	92,9	$2,9 \times 10^{-6}$
42	Mo	95,9	$8,1 \times 10^{-6}$
43	Tc	97,0	$9,5 \times 10^{-7}$
44	Ru	101,0	$3,1 \times 10^{-6}$
45	Rh	102,9	$1,6 \times 10^{-6}$
46	Pd	106,4	$4,6 \times 10^{-6}$
47	Ag	107,9	$1,5 \times 10^{-6}$
48	Cd	112,4	$4,0 \times 10^{-6}$
49	In	114,8	$8,8 \times 10^{-7}$
50	Sn	118,7	$4,7 \times 10^{-6}$
51	Sb	121,8	$9,9 \times 10^{-7}$
52	Te	127,6	$5,7 \times 10^{-6}$
53	J	126,9	$1,1 \times 10^{-6}$
54	Xe	131,3	$2,7 \times 10^{-6}$
55	Cs	132,9	$6,5 \times 10^{-7}$
56	Ba	137,3	$6,7 \times 10^{-6}$
57	La	138,9	$6,0 \times 10^{-7}$
58	Ce	140,1	$1,8 \times 10^{-6}$
59	Pr	140,9	$4,3 \times 10^{-7}$
60	Nd	144,2	$1,6 \times 10^{-6}$

Z	Particle	A_i	C_i/C_{26}
61	Pm	144,2	$1,9 \times 10^{-7}$
62	Sm	145,0	$1,8 \times 10^{-6}$
63	Eu	150,4	$3,1 \times 10^{-7}$
64	Gd	152,0	$1,4 \times 10^{-6}$
65	Tb	157,3	$3,5 \times 10^{-7}$
66	Dy	158,9	$1,4 \times 10^{-6}$
67	Ho	162,5	$5,3 \times 10^{-7}$
68	Er	164,9	$8,8 \times 10^{-7}$
69	Tu	167,3	$1,8 \times 10^{-7}$
70	Yb	168,9	$8,9 \times 10^{-7}$
71	Lu	173,0	$1,3 \times 10^{-7}$
72	Hf	175,0	$8,1 \times 10^{-7}$
73	Ta	178,5	$7,3 \times 10^{-8}$
74	W	180,9	$8,1 \times 10^{-7}$
75	Re	183,9	$2,8 \times 10^{-7}$
76	Os	186,2	$1,2 \times 10^{-6}$
77	Ir	190,2	$7,9 \times 10^{-7}$
78	Pt	192,2	$1,5 \times 10^{-6}$
79	Au	195,1	$2,8 \times 10^{-7}$
80	Hg	197,0	$4,9 \times 10^{-7}$
81	Tl	200,6	$1,5 \times 10^{-7}$
82	Pb	204,4	$1,4 \times 10^{-6}$
83	Bi	207,2	$7,3 \times 10^{-8}$
84	Po	209,0	0
85	At	210,0	0
86	Rn	222,0	0
87	Fr	223,0	0
88	Ra	226,0	0
89	Ac	227,0	0
90	Th	232,0	$8,1 \times 10^{-8}$
91	Pa	231,0	0
92	U	238,0	$4,9 \times 10^{-8}$

Not Official Standard
For Review Only

Not Official Standard
For Review Only

ICS 49.140

Price based on 6 pages

Licensed to: Tongson Nick Mr.

Downloaded: 2012-09-10

Single user licence only, copying and networking prohibited